

A persistent ~ 1 Hz quasi-periodic oscillation in the dipping low-mass X-ray binary 4U 1323–62

Peter G. Jonker¹, Michiel van der Klis¹, Rudy Wijnands¹

Subject headings: accretion, accretion disks — stars: individual (4U1323-62) — stars: neutron — X-rays: stars

ABSTRACT

We have discovered a ~ 1 Hz quasi-periodic oscillation (QPO) in the persistent-emission, the dips, and the type I X-ray bursts of the low-mass X-ray binary 4U 1323–62. The rms amplitude of the QPO is approximately 9%, only weakly depending on photon energy. The amplitude is consistent with being constant throughout the persistent-emission, the dips and the bursts in all but one observation, where it is much weaker during one dip. These properties suggest that we have observed a new type of QPO, which is caused by quasi-periodic obscuration of the central X-ray source by a structure in the accretion disk. This can only occur when the binary inclination is high, consistent with the fact that 4U 1323–62 is a dipping source. The quasi-periodic obscuration could take place by partial covering of an extended central X-ray source by a near-opaque medium, or by covering of a point source by a medium of suitable characteristics to produce the relatively energy-independent oscillations.

¹Astronomical Institute “Anton Pannekoek”, University of Amsterdam, and Center for High-Energy Astrophysics, Kruislaan 403, 1098 SJ Amsterdam; peterj@astro.uva.nl, michiel@astro.uva.nl, rudy@astro.uva.nl

1. Introduction

The low-mass X-ray binary (LMXB) 4U 1323–62 shows both type I X-ray bursts and periodic dips. The bursts and dips were discovered with the EXOSAT satellite (van der Klis et al. 1984, 1985a,b). The periodic dips in the light curves of dipping LMXBs are thought to be caused by periodic obscuration of the central source by a structure formed in an interplay between the gas stream from the companion and the accretion disk (White & Swank 1982; White, Nagase, & Parmar 1995); the period of the dips is the orbital period.

In 4U 1323–62, the dip period is 2.93 hours and the dips last ~ 50 minutes (van der Klis et al. 1985a,b; Parmar et al. 1989). From the burst properties and peak fluxes, and from the absence of eclipses the distance of the source is inferred to be 10–20 kpc, and the inclination $\leq 80^\circ$. The energy spectrum fits an absorbed power law with energy spectral index $\alpha = 0.53 \pm 0.07$, and $N_H = (4.0 \pm 0.3) \times 10^{22}$ atoms cm^{-2} .

In non-dipping LMXBs, numerous QPO phenomena have been reported, at frequencies of several Hertz to > 1200 Hz (van der Klis 1995, 1998 for reviews). So far observations of any < 100 Hz QPO phenomena in dipping LMXBs have been lacking (kHz QPOs were recently reported in 4U 1916–05 by Barret et al. 1997). In this Letter we report the discovery of ~ 1 Hz QPOs in the dipper 4U 1323–62. The frequency as well as the amplitude of the QPO is approximately constant throughout the bursts, dips, and persistent-emission. This is the first QPO reported to persist both through type I X-ray bursts and the persistent-emission, and the first < 100 Hz frequency QPO in a dipper.

2. Observations and analysis

4U 1323–62 was observed five times with the proportional counter array (PCA; Jahoda et al. 1996) on board the RXTE (Rossi X-ray Timing Explorer; Bradt, Rothschild & Swank 1993) satellite on 1997 April 25, 26, and 27 (see Table 1). The total amount of good data was 80 ksec. During each observation one or two bursts and one or two dips were observed, resulting in a total of seven type I X-ray bursts and eight dips (some of which were only partially observed). Two bursts in observation 3 showed a secondary burst ~ 500 s after the primary one. These last approximately 30 s, rise quickly up to count rates of 370 c/s (2–60 keV; this is $\sim 21\%$ of the primary burst peak count rate) and 650 c/s ($\sim 37\%$) and decrease slowly. A burst in observation 2 took place during a dip. The dips lasted ~ 60 minutes, slightly longer than those reported by Parmar et al. (1989). The mean background-subtracted persistent 2–60 keV count rate changed little between observations (see Table 1). In observation 4, during 40% of the time only 4 of the 5 PCA detectors were active. All reported count rates are for 5 detectors.

Data were obtained over an energy range of 2–60 keV with 16 s time resolution in 129 energy channels, and simultaneously with a time resolution of $1 \mu\text{s}$ in 255 energy channels.

In Figure 1 (top panel), we show part of the light curve of observation 1, showing a dip, a burst, and the persistent-emission. The dip starts ~ 600 s after the beginning of the observation and lasts until ~ 4300 s. We also show the corresponding hardness curve (lower panel); the hardness is defined as the 16-s averaged ratio of the count rate in the 9.7–16 keV energy band to that in the 2–7.9 keV

band. During the dips the hardness ratio is higher than in the persistent emission, and at the burst start the hardness ratio decreases.

Using the light and hardness curves, we divided the data into three different categories; the dips, the bursts and the persistent emission. This led to the definition of time intervals during which the source is in one of these three states. When analyzing the data the same time intervals were applied in each energy band. We defined dip ingress by the decrease in count rate and simultaneous increase in hardness evident in Figure 1; dip egress is the inverse. In between dip ingress and egress, the count rate and hardness behavior is erratic during ~ 60 minutes. Sometimes during a dip the persistent-emission count rate and hardness levels are briefly reached. We still considered those data to be part of the dip. The onset of a burst is characterized by a steep rise in the count rate simultaneously with a decrease in the hardness ratio. We determined the burst decay time by fitting an exponential to the (2–60 keV) decay at a time resolution of 1/8th second. The end of the burst was taken as three times the e-folding time after the onset. One burst decay time was 60 ± 2 s, five were consistent with 82 s and one was 92 ± 3 s. In our power spectral analysis of the bursts concerning the ~ 1 Hz QPO, we only used the burst decay interval; in our search for > 100 Hz burst oscillations we also used the burst rises.

We calculated power spectra separately for the persistent-emission and for the dips using 64-s segments with a time resolution of 1 ms in the 2–60 keV band, as well as in four energy bands (2–5.0, 5.0–6.4, 6.4–8.6, 8.6–13.0 keV). We applied a similar analysis to the bursts, but using 16-s segments enabling us to obtain 5 power spectra for most of the bursts, and reducing the low frequency component in the

power spectra due to the burst profile.

In each energy band all 1/64–512 Hz power spectra corresponding to dips (320 in number), or persistent emission (881) were averaged. The power spectra were fitted with a fit function consisting of a Lorentzian (the QPO), an exponentially cut-off power law, and a constant to represent the Poisson noise. In case of the (1/16–512 Hz) power spectra calculated from the 16-s segments (32) obtained during bursts, an extra Lorentzian centered on ~ 0 Hz was used in order to account for the power spectral component due to the burst profile. The errors on the fit parameters were determined using $\Delta\chi^2 = 1.0$ (1σ single parameter), and upper limits by using $\Delta\chi^2 = 2.71$, corresponding to a 95% confidence level.

3. Results

In the persistent-emission power spectra, we discovered a very significant (31σ) ~ 1 Hz QPO (Fig. 2). Its frequency was 0.87 ± 0.01 Hz in observation 1, 2, and 3, and 0.77 ± 0.01 Hz in observation 4 and 5. This shift of ~ 0.1 Hz between observations is not correlated to the changes in the persistent count rate level (Table 1). The QPO was detected during the persistent emission as well as during the dips and the bursts. It can be directly observed in the light curves of the bursts (see Fig. 3). During a burst the amplitude of the QPO increases by a factor of ~ 10 to keep the fractional amplitude approximately constant (see below). The Poisson counting noise prevented us to directly see the QPO in the non-burst parts of the lightcurve.

Except during the dip in observation 4 when the fractional rms amplitude was only $2.2 \pm 0.2\%$ and the FWHM only 0.07 ± 0.02 Hz, the frequency, 2–60 keV fractional rms

amplitude, and FWHM of the QPO are consistent with being the same for the dips and the persistent-emission in each observation at values of 0.77 or 0.87 Hz, 9%, and 0.25 Hz, respectively. To improve the signal to noise, we averaged the power spectra corresponding to the persistent emission and the dips across all observations in our further analysis. The FWHM and frequency of the QPO in all the persistent-emission and dip data combined is 0.25 ± 0.01 Hz and 0.84 ± 0.02 Hz, respectively; in the individual energy bands we found numbers consistent with this. In determining the fractional rms amplitudes reported below, we fixed FWHM and frequency to these values. The 2–60 keV fractional rms amplitudes in all observations combined have values of $9.1 \pm 0.1\%$, $8.9 \pm 0.4\%$, and $10.9 \pm 0.5\%$ for persistent emission, dips, and bursts, respectively. While the persistent-emission and the dip rms amplitudes are identical within the errors, the rms amplitude of the QPO during the bursts is slightly higher. Systematic errors in the rms normalization due to the trend in the count rate in the 16-s data segments caused by the burst profile, and due to the interaction in the fit procedure with the extra Lorentzian component can probably account for this small difference. We estimate their influence to be of the order of the discrepancy, since with 64-s burst power spectra, where both of these effects are more prominent, we obtained an rms amplitude of $11.9^{+3.4}_{-0.9}\%$.

The rms amplitude of the QPO only weakly depends on energy (Table 2); in the persistent-emission it is consistent with a small increase towards higher energies. The fractional rms amplitudes during the dips, and the bursts are consistent with this small increase. No time delay was found in the QPO (0.7–0.9 Hz) between the 2.8–7.5 keV and 7.5–60 keV energy bands, with a 95% confidence upper

limit of 14.5 ms for a soft lag (1.8×10^{-2} times the QPO cycle) and an upper limit of 5.4 ms (6.8×10^{-3} times the QPO cycle) for a hard lag.

We searched for kHz QPOs, but none were found with upper limits of 8–10% for a fixed FWHM of 25 Hz over a frequency range of 100–1000 Hz in all persistent-emission and dip data combined. These upper limits do not exclude the presence of kHz QPOs since many other sources have kHz QPO fractional rms amplitudes below this (van der Klis 1998).

We set upper limits on any band-limited noise component by adding an exponentially cut-off power law to the fit function. Fixing the power law index to values of 0–1, we derived upper limits of 4.2%–6.4%.

We searched the primary and secondary burst data for high frequency burst oscillations in various ways. In order to increase the sensitivity we averaged the burst rise power spectra of the different bursts using different energy bands (see also Miller 1998). No high frequency burst oscillations were found with upper limits from 0.5-s power spectra varying between 24% at the top and 45% near the end of the burst.

For reference we fitted the persistent-emission energy spectrum of part of observation 2 with a model consisting of an absorbed power law and a gaussian line at 6.5 ± 0.2 keV with a FWHM of ~ 1 keV. The fit was good, with a χ^2 per degree of freedom of 36/40. The photon index of the power law was 1.75 ± 0.02 , and the hydrogen column density is $(4.0 \pm 0.3) \times 10^{22}$ atoms cm^{-2} . This spectrum implies a 2–25 keV flux of $1.5 \times 10^{-10} \text{ erg s}^{-1}$. Using a distance of 10 kpc this results in a 2–25 keV luminosity of $1.8 \times 10^{36} \text{ erg s}^{-1}$. The spectrum is similar to that previously found by Parmar et al. (1989), although our photon index is somewhat steeper.

4. Discussion

We have discovered a QPO with a frequency of approximately 1 Hz. This QPO shows a unique combination of properties;

1) it is observed during the bursts, dips and persistent emission with the same rms amplitude (except during the dip in observation 4);

2) the rms amplitude of the QPO depends only weakly on photon energy;

3) no band-limited noise component was found with upper limits of $\sim 5\%$.

In all types of LMXBs, QPOs are known to occur with frequencies ranging from 0.01 to 1200 Hz (see van der Klis 1995, 1998 for reviews). Low frequency (0.01–10 Hz) QPOs are known to occur in a number of black hole candidates and atoll sources (see van der Klis 1995 for a review). Unlike the QPO we found in 4U 1323–62, the fractional amplitudes of low frequency QPO in these sources depend strongly on photon energy, and the QPOs are found to be superimposed on a strong band-limited noise component (Wijnands & van der Klis 1998). In the Z sources (the most luminous neutron star LMXBs) low frequency (5–20 Hz) QPOs are also well known (van der Klis 1995), the so called NBOs. However these QPOs occur at or around the Eddington mass accretion rate, which is different from the case in 4U 1323–62. Perhaps our ~ 1 Hz QPO is related to that reported by Kommers et al. (1998) in the pulsar 4U 1626–67, which they suggest is caused by a structure orbiting the neutron star. However there is no information on the energy dependence of the 0.048 Hz QPO. We conclude that so far the ~ 1 Hz QPO seems to be in a class of its own.

Models for the ~ 1 Hz QPO must explain why the rms amplitude of the QPO is constant during bursts, dips and persistent-emission. In models involving a radiation-dominated

(inner) disk, we would expect these properties to be different during a burst. Therefore, we consider explanations of the QPO in terms of modulations caused by wave packets of sound waves in the disk (Alpar & Yilmaz 1997) and in terms of fluctuations in the electron scattering optical depth at the critical Eddington mass accretion rates (Fortner et al. 1989) to be unlikely. Furthermore, assuming isotropic emission, the 2–25 keV flux in 4U 1323–62 is too low to produce these fluctuations in the way proposed by Fortner et al. (1989).

Since this QPO is observed in a dipper, we suggest an origin related to the high inclination at which we observe this system. The ~ 1 Hz QPO cannot be formed by a ripple in the same structure that is also causing the dips, since the fractional rms amplitude of the QPO should then vary with energy in the same way as the dips. This seems not to be the case, since during the persistent emission the spectrum is softer than during the dips, whereas the QPO rms amplitude increases slightly with photon energy. The fact that the QPO properties are the same in and outside the dips also makes this model unlikely. However, quasi-periodic obscuration of the central source by a structure elsewhere in the accretion flow may cause the QPO. If we assume that the QPO is caused by structures orbiting the neutron star with a Keplerian frequency of ~ 1 Hz, the orbital radius is $\sim 2 \times 10^8$ cm for a $1.4 M_{\odot}$ neutron star. A limited lifetime of the modulating structures, and/or different or shifting radii can cause the modulation of the X-rays to be quasi-periodic instead of periodic. Using the fact that no eclipses have been observed, we derive the height above the plane of the structure causing the ~ 1 Hz QPO to be $> 5 \times 10^7$ cm for a lower main sequence companion and a neutron star of $1.4 M_{\odot}$. This scale-height is difficult to explain

in view of the Shakura-Sunyaev thin α disk model. Recently Pringle (1996) discussed a model where accretion disks in LMXBs are subject to radiation driven instabilities, causing the accretion disks to warp. If these warps are responsible for the X-ray modulations as observed, the radius and structure of the warp may change on viscous time scales, which for typical parameters is on the order of days to weeks (Frank, King, & Raine 1992). Scheduled observations of this source will enable us to test this hypothesis.

Neither inverse Compton scattering nor regular Compton scattering can cause the observed X-ray modulation on its own. Both would result in a strong energy dependence of the QPO such as has been observed for kHz QPOs in, e.g., 4U 0614+09 (Méndez et al. 1997), which is inconsistent with what we observe in 4U 1323–62. However, an intermediate-temperature structure, one containing both hot and cold electrons, or one combining Compton scattering with absorption could cause the observed modulations.

A much simpler model is possible in terms of partial covering of an extended source. Quasi-periodic obscuration of part of this central source by an opaque medium, for example an orbiting bump on the disk’s surface, can be responsible for modulations with little energy dependence as observed.

This work was supported in part by the Netherlands Foundation for Research in Astronomy (ASTRON) grant 781-76-017. This research has made use of data obtained through the High Energy Astrophysics Science Archive Research Center Online Service, provided by the NASA/Goddard Space Flight Center. We would like to thank Jeroen Homan for stimulating discussions and the referee for his com-

ments and suggestions.

REFERENCES

- Alpar, M.A., Yilmaz, A., 1997, *New Astronomy* 2, 225
- Barret, D., Olive, J.F., Boirin, L., Grindlay, J.E., Bloser, P.F., Chou, Y., Swank, J.H., and Smale, A.P. 1997, *AIU Circ.* 6793
- Bradt, H.V., Rothschild, R.E., Swank, J.H., 1993, *A&AS*, 97, 355
- Fortner, B., Lamb, F.K., Miller, G.S., 1989, *Nature*, 342, 775
- Frank, J., King, A., Raine, D., 1992, *Accretion Power in Astrophysics* (Cambridge: Cambridge Univ. Press)
- Jahoda, K., Swank, J.H., Giles, A.B., Stark, M.J., Strohmayer, T., Zhang, W., Morgan, E.H., 1996, *SPIE*, 2808, 59
- Méndez, M., van der Klis, M., van Paradijs, J., Lewin, W.H.G., Lamb, F.K., Vaughan, B.A., Kuulkers, E., and Psaltis, D. 1997, *ApJLetters*, 485, 37
- Miller, M.C., 1998, *ApJ*, submitted, *astro-ph/9809235*
- Parmar, A.N., Gottwald, M., van der Klis, M., & van Paradijs, J., 1989, *ApJ*, 338, 1024
- Pringle, J.E., 1996, *MNRAS*, 281, 357
- van der Klis, M., van Paradijs, J., Jansen, F.A., & Lewin, W.H.G., 1984, *IAU Circ.* 3961
- van der Klis, M., Parmar, A.N., van Paradijs, J., Jansen, F.A., & Lewin, W.H.G., 1985a, *IAU Circ.*, No 4044

- van der Klis, M., et al., 1985b, Sp. Sci. Rev. 40, 287
- van der Klis, M., 1995, in X-Ray Binaries, ed. W.H.G. Lewin, J. van Paradijs, & E.P.J. van der Heuvel (Cambridge: Cambridge Univ. Press)
- van der Klis, M. 1998, in 'The many faces of Neutron Stars', R. Buccheri, J. van Paradijs & M.A. Alpar (Eds), NATO ASI Series C, Vol. 515, pp. 337-368 (Kluwer Academic Publishers)
- White, N.E., and Swank, J.H., 1982, ApJ Letters, 253, L61
- White, N.E., Nagase, F., Parmar, A.N. 1995, in X-Ray Binaries, ed. W.H.G. Lewin, J. van Paradijs, & E.P.J. van der Heuvel (Cambridge: Cambridge Univ. Press)
- Wijnands, R., van der Klis, M. 1998, ApJ in press

Fig. 1.— Top: The 2–60 keV, 16-s averaged, background-subtracted light curve beginning at the start of the first observation (see Table 1) showing a dip and a burst. No dead time corrections have been applied. The statistical uncertainty is typically 5 c/s. Bottom: The 16 s averaged hardness curve (9.7–16 keV/2–7.9 keV, see text) of the same data. Typical errors in the hardness are 0.03. The arrows indicate the start and end of the dip in both the light- and hardness curve.

Fig. 2.— Power density spectrum of the full (2–60 keV) energy band of the persistent emission of all the observations combined showing the ~ 1 Hz peak. The line drawn through the data points represents the best fit to the data. No dead time corrections have been applied.

Fig. 3.— Light curve showing the QPO during a burst. Five points correspond to 1.25 s, close to the QPO period.

TABLE 1

LOG OF THE OBSERVATIONS. THE AVERAGE PERSISTENT-EMISSION COUNT RATE OF EACH OBSERVATION IS GIVEN IN COLUMN 5.

Number	Observation	Date	Time (UTC)	Persistent emission 2–60 keV (c/s/5PCU)
1	20066-02-01-00	25 April 1997	22:03–03:56	106
2	20066-02-01-03	26 April 1997	05:06–09:41	104
3	20066-02-01-01	26 April 1997	22:02–03:55	95
4	20066-02-01-04	27 April 1997	05:08–09:42	102
5	20066-02-01-02	27 April 1997	23:12–03:54	94

TABLE 2

FRACTIONAL RMS AMPLITUDES (IN %) OF THE ~ 1 Hz QPO. IN DETERMINING THESE VALUES THE FWHM AND FREQUENCY WERE FIXED TO 0.25 AND 0.84 Hz, RESPECTIVELY.

Category	Energy (keV)				2–60
	2–5.0	5.0–6.4	6.4–8.6	8.6–13.0	
Persistent	8.0 ± 0.3	9.9 ± 0.5	9.8 ± 0.4	10.9 ± 0.4	9.1 ± 0.1
Dips	8.1 ± 0.9	9.2 ± 1.1	10.2 ± 0.8	9.7 ± 0.8	8.9 ± 0.4
Bursts	10.3 ± 0.9	10.2 ± 0.9	10.0 ± 1.0	10.6 ± 1.0	10.9 ± 0.5





